

GENERAL INFORMATION

RESIST II™ contains potassium, which is intended to stimulate root development and resistance to disease.

RESIST II™ is non-phytotoxic when used as directed and is compatible with most commonly used pesticides and fertilizers.

RESIST II™ has a low salt index and can be applied as a foliar or used in drip or micro-sprinkler irrigation systems.

RESIST II™ should not be applied to plants that are stressed by high temperature or lack of moisture.

Phosphorous acid products are intended to be used as a supplemental fertilizer treatment.

* When using **RESIST II™** as a foliar application, the phosphite ions are taken up directly by the plant foliage and may undergo a degree of conversion to phosphate ions, or are utilized directly by plants, as phosphite ions. When using **RESIST II™** as a soil application on annual crops, you may experience a lesser response from the initial crop, with a corresponding superior response from succeeding crops. In addition, placement close to seeds or root zones may be injurious to crops. The effect may be aggravated by a soil pH below 6.5.

WARRANTY: Since weather, crops, soil and other conditions may vary, neither ACTAGRO, LLC, nor the Seller make a warranty of any kind, expressed or implied, concerning the use of this product. The user assumes all risk of use and handling whether or not in accordance with directions or suggestions.

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED. AVOID CONTACT WITH SKIN, EYES OR CLOTHING. In case of contact immediately flush skin, eyes or clothing with plenty of water for at least 15 minutes.

CAUTION: If any other chemicals are to be used in the mixture, be sure to check compatibility. Do not mix with copper based products.



GUARANTEED ANALYSIS

Total Phosphoric Acid (P₂O₅) 27.0%
Soluble Potash (K₂O)..... 25.0%

Derived from potassium hydroxide and phosphorous acid.

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Net Contents 5 gallons (18.9 liters)

12.1 lbs per gallon (1474 grams/liter) @ 68°F

DIRECTIONS FOR USE

RESIST II™ can be applied to most vegetable and cucurbit crops, field crops, row crops, citrus, fruit and nut trees, vines, olives, turf, nursery and greenhouse plants and ornamental farms.

RESIST II™ may be applied to the soil or used as a foliar and may be applied through center pivot irrigation systems.

SOIL APPLICATION RATES

ALL CROPS: Apply to the soil at 1 quart to 2 quarts per acre. **RESIST II™** is most effective with drip or micro-sprinkler irrigation systems. Repeat as needed.

ORNAMENTAL -NURSERY-GREENHOUSE: Apply 1/2 oz to 1 oz per 1000 sq ft. Repeat as needed.

FOLIAR APPLICATION RATES

RESIST II™ may be applied by air or ground spray. Use the appropriate volume of water for the application method and the crop. Apply 1 quart to 2 quarts per acre. Repeat as needed.

Application Precaution: As with all acid materials, user should exercise precaution to avoid potential foliar burning. Use caution if applying this product in combination with other materials since mixing could result in a potentially hazardous reaction caused by product incompatibilities. A compatibility test is recommended before tank mixing with other products. The pH of any final spray solution should be checked and the appropriate amount of the solution should be applied to a small area of the intended crop to ensure plant compatibility prior to the application. **THE USER ASSUMES FULL RESPONSIBILITY TO ENSURE PRODUCT, EQUIPMENT AND PLANT COMPATIBILITIES.**

Mixing Instructions

RESIST II™ mixes easily with water as follows:

1. Partially fill tank with water and start agitation.
2. Add pesticides or other nutrients.
3. Add **RESIST II™** to circulating tank.
4. Fill tank and agitate thoroughly.

California

ACTAGRO, LLC.
PO BOX 309 • BIOLA, CA 93606
Copyright 2006